

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/18126**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(7) :A61K 49/00

US CL :424/9.1

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/1.11, 1.65, 9.1, 9.2, 9.3, 9.4, 9.5, 9.6, 9.7, 9.8

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

BIOSIS MEDLINE USPATFULL EMBASE CAPLUS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5,618,831 A (SHISHIDO et al) 08 April 1997, see entire document, especially, abstract; column 2, lines 22-60.	1-3
Y	IYPE et al. Carcinogenesis by Nonmutagenic Chemicals: Early Response of Rat Liver Cells Induced by Methapyrilene. Cancer Research. May 1985, Vol. 45, pages 2184-2191, especially, abstract; page 2185, column 1, third complete paragraph; page 2185, columns 1-2, bridging paragraph; page 2186, column 1, third paragraph.	1-3

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

29 AUGUST 2000

Date of mailing of the international search report

07 SEP 2000

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

D. Jones

Telephone No. (703) 308-1235

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
10 January 2002 (10.01.2002)

PCT

(10) International Publication Number
WO 02/02149 A1

- (51) International Patent Classification⁷: **A61K 49/00**
- (21) International Application Number: **PCT/US00/18126**
- (22) International Filing Date: **30 June 2000 (30.06.2000)**
- (25) Filing Language: **English**
- (26) Publication Language: **English**
- (71) Applicant (*for all designated States except US*): **ZILA, INC.** [US/US]; 5227 North Seventh Street, Phoenix, AZ 85014-2800 (US).
- (72) Inventor; and
- (75) Inventor/Applicant (*for US only*): **BURKETT, Douglas, D.** [US/US]; 4736 E. Euclid, Phoenix, AZ 85044 (US).
- (74) Agent: **DRUMMOND, William, H.**; Drummond & Duckworth, 4590 MacArthur Boulevard, Suite 500, Newport Beach, CA 92660 (US).
- (81) Designated States (*national*): AT, AU, BR, CA, CN, CZ, HU, IL, IN, JP, KR, MX, NO, NZ, PL, RO, SG, SK, TR, UA, US, ZA.
- (84) Designated States (*regional*): Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).
- Published:**
— *with international search report*
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*



WO 02/02149 A1

(54) Title: **RHODAMINE DIAGNOSTIC AGENT AND DIAGNOSTIC METHODS FOR DETECTION OF EPITHELIAL CANCER**

(57) Abstract: Rhodamine dye compositions and methods for detecting and/or delineating cancerous and precancerous epithelial tissue.

-1-

**RHODAMINE DIAGNOSTIC AGENT
AND DIAGNOSTIC METHODS FOR DETECTION
OF EPITHELIAL CANCER**

This invention relates to a novel diagnostic agent for detection of cancerous and precancerous epithelial tissue.

According to another aspect, the invention pertains to novel methods for detecting and/or delineating cancerous and/or precancerous tissue of the epithelium.

In another respect the invention concerns to such diagnostic procedures and agents useful therein which are especially useful for in vivo screening of patients for possible oral cancer as part of routine dentist's or physician's examinations or procedures, such as periodic dental or physical examinations, dental cleaning, etc.

In yet another aspect the invention relates to such procedures and compositions useful therein, which use dye stains that are more readily available and/or less expensive and less complicated to synthesize and/or purify than the dyes employed in prior art procedures.

-2-

In yet another respect, the invention concerns such in vivo procedures and compositions employing a dye which, despite prior art teachings otherwise, is sufficiently non-toxic that it can be employed by rinsing the entire oral cavity and/or gargling.

In-vivo diagnostic procedures for detecting premalignant epithelial lesions, such as oral lesions and oral carcinomas, employing dye compositions that are selectively retained by tissues rendered abnormal by dysplasia, hyperplasia, tumorigenesis and other active surface lesions, are known in the art. For example, procedures employing fluorescein or fluorescein derivatives are disclosed in Chenz, Chinese Journal of Stomatology (27:44-47(1992)) and Filurin (Stomatologia (Russian) 72:44-47 (1993)). These procedures involve application of the dye, followed by visual examination under ultraviolet light to detect cancerous/precancerous tissue, which is selectively fluorescent.

Another prior art procedure involves in vivo application by rinsing with toluidine blue O, followed by normal visual examination to detect any selectively

-3-

stained tissue. Such procedures are disclosed, for example, in U.S. Patent 5,372,801 to Tucci, et al. and in U.S. Patent 4,321,251 to Mashberg. Toluidine blue has been used for decades as a histopathological stain for in-vitro use. Through this use it has become known as a metachromatic dye, staining nuclei rich in DNA and RNA a purple to pink color. The inherent deep blue color of toluidine blue O is changed to purple or pink when the dye is bound to nucleic acid or other acidic cellular macromolecules. Of course, this type of staining is dependent on the dye gaining access to internal subcellular structures such as the nucleus. Such access is readily obtained only by "fixing" a tissue sample with formaldehyde or other reagent that disrupts the cellular membrane without destroying general cellular structure.

In contrast to the mechanism involved in in-vitro use, the staining of oral tissue in vivo by toluidine blue O is due to its ability to penetrate the cell walls and attach to the mitochondria, which retains the dye longer than components of the extracellular matrix.

-4-

The Mashberg procedure involves application of the toluidine blue O solution as a rinse of the entire oral cavity, with gargling, followed by rinses with water and acetic acid to remove dye that is not retained by the cancerous or precancerous tissue. The preliminary diagnosis by the Mashberg procedure is then confirmed by direct application of the toluidine blue O composition to the suspect site 10-14 days later. The Tucci '801 patent discloses an improved toluidine blue O composition for use according to the general procedure taught by Mashberg.

An in vivo procedure involving use of Lugol's solution (iodine) and toluidine blue O was proposed for detecting esophageal cancer synchronous with upper aerodigestive tract cancers in Papazian, Gastroenterologic Clinique et Biologique 9:16-22 (1985).

More recently, Pomerantz U.S. Patent 5,882,627 disclosed a structurally defined class of oxazine and thiazine dyes that are useful in general accordance with the Mashberg diagnostic protocol.

-5-

Bernal et al., Cancer Research 43, 716-720 (1983) disclosed that rhodamine dye selectively inhibits the growth of and kills carcinoma cells in vitro. Gaboury et al., U.S. Patent 5,773,460 discloses that rhodamine is preferentially retained by many tumor cells and proposes that certain esters of rhodamine are useful for vitro photodynamic inhibition of certain tumor cell lines, but indicates that systemic toxicity may limit its usefulness in chemotherapy.

Rhodamine , (2-(6-amino-3-imino-3H-xanthen-9-yl) benzoic acid methyl ester and ionic salts thereof, e.g., hydrochloride salts, is a lipophilic cationic dye of the pyrylium class. It is effective to selectively identify and/or delineate cancerous and precancerous epithelial tissue by topical application to the epithelium, followed by normal visual examination, in general accordance with the protocol disclosed by the Mashberg '251 patent.

Suitable compositions of rhodamine for application of the dye to epithelial tissue are prepared by mixing the dye with a suitable pharmaceutically acceptable solvent. Preferably the pH of the rhodamine solution is

-6-

adjusted with a suitable pharmaceutically acceptable buffer system to yield a final solution that is substantially isotonic and has a pH in the range of approximately 2.5 to 7.0, preferably 4.0-5.0. This can be accomplished by an acetic acid-sodium acetate buffer system. Other suitable buffer systems include citric acid-sodium citrate or mixed acid salt systems such as citric acid-sodium phosphate and the like.

The solvent used to provide the liquid rhodamine dye compositions of the invention is an aqueous solvent. According to the presently preferred embodiment of the invention, the solvent included a pharmaceutically acceptable, i.e., non-toxic, non-reactive alcohol, e.g., ethyl alcohol. Such solvents do not appreciably interfere with the staining mechanism and do not themselves contribute to the reduction of chromo forms of the dye to leuco forms.

Flavoring, stable to the other components of the dye composition, may be added to improve the palatability of the composition if it is to be used as an oral "rinse."

-7-

The amount of rhodamine dye in the liquid composition is preferably adjusted to yield a concentration of approximately 1% by weight of the final composition, although higher concentrations can be employed and lower concentrations are at least partially effective. At present, I prefer to employ dye compositions containing from about 0.5 to about 3.5% by weight of the rhodamine component.

The invention also contemplates compositions for use in accordance with the methods of the invention, in which any leuco form of the dye present in the composition is oxidized to the chromo form, by inclusion of a pharmaceutically acceptable oxidizing agent, in the manner analogous to that disclosed in the Tucci '801 patent.

EXAMPLES

The following examples are presented in order to illustrate practice of the invention to those skilled in the art and not by way of limitation of the scope thereof, which is defined only by the appended claims.

-8-

Example 1**Preparation of Diagnostic Composition**

A diagnostic composition is prepared by mixing each of the indicated components in the following proportions (% by weight):

Purified Water U.S.P.	83.85
Glacial Acetic Acid U.S.P.	4.61
Sodium Acetate Trihydrate U.S.P.	2.45
SD18 Ethyl Alcohol	7.48
Hydrogen Peroxide 30%, U.S.P.	0.41
IFF Raspberry IC563457	0.20
Rhodamine	1.00

Example 2**Preparation of Rinse Solution**

A rinse solution is prepared by mixing the following components in the indicated proportions (weight %):

-9-

Purified Water U.S.P.	98.70
Glacial Acetic Acid U.S.P.	1.00
Sodium Benzoate U.S.P.	0.10
IFF Raspberry IC563457	0.20

Example 3

Clinical Effectiveness

The clinical effectiveness of the compositions of Examples 1 and 2, is compared to toluidine blue O diagnostic control compositions prepared in accordance with the Tucci '801 patent, using the diagnostic protocol disclosed in the Mashberg '251 patent.

Patients are first screened for oral pathology employing the TBO control composition. After identifying potential cancerous or precancerous pathology, all traces of the TBO are removed by repeatedly rinsing the suspect sites with water and the acetic acid rinse of Example 2.

Those patients exhibiting oral pathology are then used as test subjects for the rhodamine diagnostic

-10-

composition of Example 1. 2-3 cc of the rhodamine composition is applied by painting the pathologic mucosal surface, followed by rinsing with the rinse mixture and water to remove excess rhodamine composition.

Histological examination of tissue from the areas stained by the rhodamine diagnostic composition of Example 1 confirms that the rhodamine composition is at least as effective as toluidine blue O in identifying and delineating cancerous and precancerous epithelial tissue.

Example 4

The procedures of Example 3 are repeated, except that the test and control compositions are applied to the oral mucosa by rinsing, with gargling, instead of by direct application to the locus of the suspect sites. Equivalent results are obtained.

Having disclosed my invention in such terms as to enable those skilled in the art to understand and practice it, and having disclosed the presently preferred embodiment, I CLAIM:

claims

-11-

1. A diagnostic method for identifying and delineating cancerous epithelial tissue, comprising:

- (a) applying rhodamine to the epithelium, to selectively mark cancerous tissue; and
- (b) visually examining said epithelium, to identify and delineate suspected cancerous tissue sites, by comparing the color of the suspected sites with the color of the adjacent tissue.

-12-

2. In a method for in vivo detection of premalignant epithelial lesions and carcinomas, including the steps of sequentially

rinsing the epithelium with dye stain composition which is selectively retained by cancerous and precancerous tissue, wherein the stain composition consists essentially of toluidine blue O, and

rinsing the epithelium with a rinse composition for removing unretained stain composition,

the improvement in which the stain composition comprises rhodamine.

-13-

3. A biological stain composition for in vivo detection of cancerous and precancerous tissue, comprising

- (a) rhodamine,
- (b) a pharmaceutically acceptable aqueous solvent, and
- (c) a pharmaceutically acceptable oxidizing agent for leuco rhodamine.